



Published in final edited form as:

*Arch Intern Med.* 2010 February 22; 170(4): 381–388. doi:10.1001/archinternmed.2009.533.

## Getting to “No”:

### Strategies Primary Care Physicians Use to Deny Patient Requests

**Dr. Debora A. Paterniti, PhD, Dr. Tonya L. Fancher, MD, MPH, Ms. Camille S. Cipri, BS, Dr. Stefan Timmermans, PhD, Dr. John Heritage, PhD, and Dr. Richard L. Kravitz, MD, MSPH**  
Center for Healthcare Policy and Research (Drs Paterniti and Kravitz and Ms Cipri), Division of General Medicine (Drs Paterniti, Fancher, and Kravitz), and Department of Sociology (Dr Paterniti), University of California, Davis, Sacramento; and Department of Sociology, University of California, Los Angeles (Drs Timmermans and Heritage)

### Abstract

**Background**—Physicians need strategies for addressing patient requests for medically inappropriate tests and treatments. We examined communication processes that physicians use to deal with patient requests of questionable appropriateness.

**Methods**—Data come from audio-recorded visits and postvisit questionnaires of standardized patient visits to primary care offices in Sacramento and San Francisco, California, and Rochester, New York, from May 2003 to May 2004. Investigators performed an iterative review of visit transcripts in which patients requested, but did not receive, an antidepressant prescription. Measurements include qualitative analysis of strategies for communicating request denial. The relationship between strategies and satisfaction reports in postvisit questionnaires was examined using the Fisher exact test.

**Results**—Standardized patients requested antidepressants in 199 visits; the antidepressants were not prescribed in 88 visits (44%), 84 of which were available for analysis. In 53 of 84 visits (63%), physicians used 1 or more of the following 3 strategies that explicitly incorporated the patient perspective: (1) exploring the context of the request, (2) referring to a mental health professional, and (3) offering an alternative diagnosis. Twenty-six visits (31%) involved emphasis on biomedical approaches: prescribing a sleep aid or ordering a diagnostic workup. In 5 visits (6%),

---

©2010 American Medical Association. All rights reserved.

Correspondence: Debora A. Paterniti, PhD, Division of General Medicine, University of California, Davis, 4150 V St, Ste 2500, Sacramento, CA 95817 (dapaterniti@ucdavis.edu).

**Financial Disclosure:** Dr Kravitz has received unrestricted research grants from Pfizer during the past 3 years.

**Previous Presentation:** This study was presented in part as an oral presentation at the 2009 Society of General Internal Medicine National Meeting; May 13–16, 2009; Miami, Florida.

**Additional Contributions:** The authors thank Blue Shield of California; the University of California, Davis, Primary Care Network; Western Health Advantage; Kaiser Permanente; Brown & Toland IPA; and Excellus BlueCross BlueShield. We are also indebted to 18 SPs; to participating physicians and their office staff members; to Nolan Zane, PhD (Asian American Center for Disparities Research), for data analysis and manuscript preparation; and to Mark Henderson, MD, for his thoughtful comments on an earlier draft.

**Author Contributions:** Drs Paterniti and Fancher contributed equally to the creation of this manuscript—the order of their names in the byline is entirely arbitrary. *Study concept and design:* Paterniti, Fancher, and Kravitz. *Acquisition of data:* Kravitz. *Analysis and interpretation of data:* Paterniti, Fancher, Cipri, Timmermans, Heritage, and Kravitz. *Drafting of the manuscript:* Paterniti, Fancher, and Kravitz. *Critical revision of the manuscript for important intellectual content:* Paterniti, Cipri, Timmermans, Heritage, and Kravitz. *Statistical analysis:* Kravitz. *Obtained funding:* Kravitz. *Administrative, technical, and material support:* Paterniti, Fancher, Cipri, Timmermans, Heritage, and Kravitz. *Study supervision:* Paterniti and Kravitz.

physicians rejected the request outright. Standardized patients reported significantly higher visit satisfaction when approaches relying on the patient perspective were used to deny the request ( $P=0.001$ ).

**Conclusions**—Strategies for saying no may be used to communicate appropriate care plans, to reduce provision of medically inappropriate services, and to preserve the physician-patient relationship. These findings should be considered in the context of physician education and training in light of increasing health care costs.

Patient requests for clinical care, including requests for medication, can influence physicians' decisions about treatment.<sup>1</sup> Patients make a request for medication in roughly 10% of office visits, and most requests are honored.<sup>2,3</sup> Medications prescribed at the behest of patients may not always represent physicians' first choice of treatment, particularly if the requests are commercially motivated, as for example, by direct-to-consumer advertising (DTCA).<sup>2,4,5</sup> Nevertheless, physicians are cautious when rejecting patient requests for services, in part because of physicians' perception that rejection may lower patient satisfaction.<sup>6,7</sup> Yet, data from patients are conflicting: nonfulfillment is associated with lower patient satisfaction in some studies<sup>7-10</sup> but not in others.<sup>11</sup> It may be that patient satisfaction has more to do with what the physician says or does not say than whether or not an expected prescription is offered.<sup>12</sup>

In the popular business book, *Getting to Yes: Negotiating Agreement Without Giving In*, Fisher et al<sup>13</sup> present general strategies for bridging gaps and achieving compromise in everyday life. In the clinical setting, however, "getting to yes" is not always appropriate for at least 3 reasons. First, on ethical grounds, physicians have a duty to avoid doing harm and to maximize patient benefit. This duty may conflict with other ethical obligations, such as respecting patient autonomy. Nevertheless, bioethicists are nearly unanimous that physicians are not obligated to provide unnecessary or inappropriate care.<sup>14</sup> Second, from a safety perspective, physicians must exercise caution when prescribing new, poorly tested, or marginally indicated medications.<sup>15</sup> Third, from a policy perspective, achieving control of health care costs is a critical national priority.<sup>16</sup> Physicians must balance the needs of individuals with those of society; in some cases, serving as social stewards might mean forgoing otherwise clinically indicated but very expensive care. Therefore, judicious clinical restraint in the patient visit may benefit the patient and will be a cornerstone of any effective cost-containment program.

Nevertheless, "getting to no" is not easy, and, to our knowledge, there are no studies directly examining the approaches that physicians use in everyday practice. This article examines the conversational and clinical rejection strategies that physicians use in everyday practice to deal with patient requests that they do not wish to fulfill.

## METHODS

### DESIGN

To examine pathways to rejection and to identify strategies that will allow physicians to maintain control of the treatment plan while potentially preserving patient satisfaction, we analyzed data from a randomized trial on the prescribing behavior of primary care

physicians in response to standardized patient (SP) requests for antidepressant medication.<sup>1</sup> Data included transcripts from office visits in primary care practices and postvisit questionnaires with measures of SP-reported visit satisfaction. The SPs were scripted and trained to portray 2 different clinical roles (major depression with wrist pain or adjustment disorder with back pain) while making 1 of 3 different requests (brand-specific antidepressant medication request, general request for antidepressant medication, or no request).<sup>1</sup> Investigators told SPs that they were interested in an array of physician responses to the different clinical roles and request types. The SPs who were enacting the 2 roles were trained separately and portrayed a single role for the entire study. Although the SPs were not blinded to the study design, they were not aware of specific study hypotheses. Those SPs who were instructed to make an antidepressant request were asked to make their initial request within the first 10 minutes of the visit or before the physical examination and to make a second request during the office visit if the first request did not lead to a prescription for an antidepressant.

### SETTING, PATIENTS, AND INTERVENTION

Data for the randomized trial were collected between May 2003 and May 2004. A total of 152 primary care physicians consented to participate in a study using unannounced SPs to evaluate social influences on practice and competing demands in primary care. Internists and family physicians were recruited through 4 physician groups: University of California, Davis, Primary Care Network and Kaiser-Permanente in Sacramento, California; Brown and Toland Medical Group in San Francisco, California; and Excellus BlueCross BlueShield in Rochester, New York. Cooperation rates by site ranged from 53% to 61%.

Eighteen insured, middle-aged, white, female SPs were trained and randomly assigned to make 298 unannounced visits, so that most physicians enrolled in the study saw 1 patient with depression and 1 with adjustment disorder. The SPs scheduled new visits to physicians and presented with subacute fatigue and insomnia accompanied by an unrelated orthopedic complaint referable to low back strain or carpal tunnel syndrome. Details on their training and detection rates are described elsewhere.<sup>17</sup> Visits were digitally recorded using a concealed recorder; recordings were transcribed verbatim for analysis.

A summary of the trial study results is presented in the Table. Findings related to prescribing behaviors and request types,<sup>1</sup> shared decision-making behaviors,<sup>18,19</sup> physician self-reflection and rationale related to prescribing behaviors,<sup>20</sup> and exploration of suicide<sup>21</sup> are provided in detail elsewhere.

### QUALITATIVE DATA ANALYSIS

Visit transcripts were inductively reviewed and assessed for important visit components (information gathering about the physical complaint, depression-related symptoms, patient perspective related to complaint; inquiry into the nature of the advertisement or the context of the patients' complaints; information giving about depression, antidepressants, or sleep medications; presentation of a tentative diagnosis; and discussion of a treatment plan, including patient understanding, subsequent follow-up, and the possibility for prescribing an anti-depressant). The order of these components and their relationship to presenting

symptoms and request type (brand-specific or general antidepressants) were also noted. Patient requests and physician responses were abstracted from the transcripts, and a qualitative content analysis of physician responses was performed. The content analysis included development of an exhaustive list of how physicians went about denying patient requests. This list of approaches for denying requests was systematically reviewed and categorized into 3 strategic pathways to no, which were analyzed by a medical sociologist (D.A.P.) and a physician (T.L.F.) using a systematic and iterative approach to content analysis. Patterns and themes were further reviewed by all coauthors, and a final set of approaches and strategies was established by consensus. The 3 pathways to no and substrategies are outlined in Figure 1 and detailed in the “Results” section. Reviewers coded the transcripts blinded to outcome measures of patient satisfaction, to which each approach was later correlated in the analysis.

## QUANTITATIVE MEASURES AND ANALYSIS

Previous work indicates that request nonfulfillment diminishes patient satisfaction, that patient-centered communication enhances it, and that SP satisfaction is correlated with the satisfaction of real patients seeing the same physician.<sup>1,22</sup> We therefore hypothesized that certain forms of request denial would be associated with lower SP-reported satisfaction. We anticipated that an approach to request denial that incorporated aspects of the patient’s interpretation of the chief complaint (“feeling tired”) could result in a preserved relationship between the physician and the patient and therefore in higher reports of SP satisfaction.

We investigated whether there was any relationship between postvisit SP satisfaction and 1 of 3 decision pathways (patient perspective based, biomedically based, or outright rejection). In the original study, the SPs completed two, 5-point Likert-scaled items for physician satisfaction: “Thinking about the visit you just made, how would you rate the physician in terms of your overall satisfaction with care [1, excellent; 5, poor]?” “Would you want this physician for your own personal physician [1, yes, definitely; 5, no, definitely not]?” The sum of these 2 items produced a reliable scale (mean [SD], 7.12 [2.30]; range, 2–10;  $\alpha=0.90$ ) that was skewed strongly positive. We therefore split the sample near the 75th percentile to produce a dichotomous variable indicating excellent satisfaction (scale score, 9 or 10) vs less than excellent satisfaction (scale score, <9). The relationship between pathways to no and excellent satisfaction was examined using the Fisher exact test, as implemented in Stata version 10.0 (Stata Corp, College Station, Texas).

## RESULTS

### PHYSICIANS AND PRACTICES

A request for medication was made in 199 (68%) of the office visits; in 88 (44%) of those visits, the request was denied. Four of the 88 visits were only partially transcribed or unavailable for transcription owing to poor recording quality, leaving 84 visits. Of the 84 visits, 54 were to general internists and 30 were to family physicians; 59 were to male physicians and 25 were to female physicians. The age, sex, and specialty distributions of the 84 visits in which requests were denied were similar to those of the other visits ( $P>.40$  in all cases).

## GENERAL CONTENT OF RESPONSES TO PATIENTS

Each visit opened with the same chief complaint of “feeling tired” plus a physical complaint of either wrist pain (presented with symptoms of major depression) or low back pain (presented with symptoms of adjustment disorder). Physician review of both chief complaints occurred in 96% (81) of all visits. Physicians’ statements about antidepressants after SP requests included comments emphasizing the problems with antidepressant use (ie, costs, delayed onset of benefit, long-term adherence requirements, and lack of efficacy for “feeling tired” or for problems of “mild,” “situational,” or “short-term” depression) and overall reluctance to prescribe antidepressants (“I’m not a pill doctor”; “I just think they [antidepressants] are overused.”)

## APPROACHES TO GETTING TO NO

Physicians used 3 strategic pathways for denying patients’ requests for antidepressants: patient perspective-based strategies (63%), biomedically based strategies (31%), or outright rejection (6%). Figure 1 illustrates the 3 approaches, which are detailed below. Specific examples from visit transcripts of the content and how physicians said no are provided in Figure 2.

**Patient Perspective-Based Approaches**—In 53 visits (63% of the 84 total visits), physicians gathered additional data about the request and its origin and offered information tailored to the patient’s presentation of information. Three approaches emphasizing the patient’s perspective on “feeling tired” or about the rationale for requesting antidepressants included (1) exploring the context of the request, (2) seeking the advice of a counselor or mental health specialist, and (3) offering an alternative diagnosis to major depression. These approaches presume an implicit validation of depression as the appropriate diagnosis and maintain the patient’s interpretation and perspective at the core of the physician response. The most frequent of the 3 approaches, exploring the context of the request, occurred in 34 of the 84 visits (40%). Physicians’ attempts to understand the original context of the request (eg, “Where did you see the ad?” “What about the ad rang true for you?”) and inquiries about recent events leading to the visit were often followed by a negotiated timeline for addressing the patient’s symptoms, some including the possibility of prescribing an antidepressant at a later date.

Referral to a counselor or mental health professional occurred in 10 of the 84 visits (12%). Eight (or 80%) of these 10 referrals came from physicians in a health maintenance organization. Physician justifications for referral included having the patient consult with someone who could “go over things” and “make a recommendation [to the physician] about the appropriateness of medication” along with the benefit of seeing someone who might provide ways to deal with stress through “skills not pills.” Physicians provided extensive information about reasons for suggesting counseling and frequently told the patient that the referral was an opportunity for her to “talk things out with someone.”

A third strategy that made use of patient perspectives included rejection of the request for an antidepressant by offering an alternative diagnosis of “situational” or “mild” depression as the reason for the patient’s chief complaint (9 visits, or 11% of the 84 total visits). In all but

1 of these 9 visits, the SP portrayed a patient with an adjustment disorder. Physicians typically followed the alternative diagnosis with specific reasons for rejecting the patient's request, including discussing the symptoms of major depression and reiterating contextual factors described by the patient to support the alternative diagnosis.

**Biomedically Based Approaches**—In 26 visits (31% of the 84 total visits), physicians used 1 of 2 biomedically based approaches to justify rejecting the request: prescribing a sleep aid (often a sedative-hypnotic, sometimes trazodone or a low-dose tricyclic agent) or ordering a diagnostic workup to rule out alternative medical illness. In the first approach, 15 physicians (18% of visits) prescribed a sleep aid, sometimes with a sleep hygiene handout, to address the patient's chief complaint of feeling tired. During these visits, physicians emphasized the ineffectiveness of antidepressants or provided justification of treatment with sleep aids over antidepressants for fatigue. Physicians instructed the patient to “try the sleep aid” and “see how you respond to it.” Some physicians even remarked that they were giving the patient an “old-fashioned antidepressant” (ie, a low dose of trazodone). Frequently, a physician would claim that fatigue might be related to sleep disturbance caused by the musculoskeletal pain, addressing both of the patient's complaints simultaneously.

In the second approach, physicians ordered diagnostic tests to rule out thyroid disease, anemia, menopause, or diabetes. Physicians frequently acknowledged the patient request as having some merit (eg, “That's what I was thinking” “That's a possibility”) and then suggested that the request might be fulfilled if the patient first followed the physician's plan. A diagnostic workup implicitly presumes that a physical condition is to blame for the symptoms and may be, from the patient's perspective, a lesser validation of the request. Overall, physicians ordered 1 or more laboratory tests in 68 of 84 visits (81%). Physicians described plans for a diagnostic workup as the primary reason for delaying attention to a patient request for an antidepressant in 11 of the 84 encounters (13%).

**Outright Rejection Approach**—In 5 of the 84 visits (6%), physicians rejected patient requests without explanation and quickly shifted the topic to investigation of the patient's musculoskeletal complaint (eg, “Let's go through and do an examination”; “What about this low back pain?”) or further exploration of the patient history unrelated to depression history or its context. Interestingly, all 5 patient requests that generated these physician responses were general requests (eg, “Do you think medication would help me?”), not brand-specific requests, for an antidepressant.

## THE RELATIONSHIP BETWEEN GETTING TO NO AND PATIENT SATISFACTION WITH PHYSICIAN AND VISIT

The relationship between approaches to no and excellent visit satisfaction was examined using the Fisher exact test. The 26 visits with scores of 9 or 10 were classified as excellent satisfaction, and the remaining 58 with scores of less than 9 were classified as less than excellent satisfaction. The SPs were significantly more likely to report excellent visit satisfaction with approaches involving the patient perspective-based strategy (Figure 3). When the approaches were dichotomized into patient perspective-based and other strategies (combining the 5 outright rejection visits with the 3 biomedically based approaches), the SPs

reported excellent visit satisfaction in 43% of the visits in which patient perspective-based approaches were used and in 10% of the visits in which other approaches were used ( $P=.001$ ).

## COMMENT

Physicians cannot always fulfill patient requests. However, little is known about the approaches that physicians use to issue denials. In this qualitative analysis of 84 office visits, physicians used 6 approaches for denying requests for antidepressants. These approaches for getting to no were classified as patient perspective based, biomedically based, or outright rejection based on the primary reason that the physician provided for denying the patient's request. The SPs reported significantly higher visit satisfaction when the physician used a patient perspective-based strategy to deny their request for antidepressants ( $P=.001$ ).

Unfulfilled requests may have consequences for the physician-patient relationship, and physicians must learn to manage these requests in a respectful and clinically sensible fashion. Unfulfilled requests have been associated with reduced satisfaction in some studies<sup>23</sup> but not in others.<sup>24–26</sup> A vignette-based study by Shah et al,<sup>27</sup> in which a patient was denied a DTCA-based request, showed evaluations of care to be significantly associated with physician communication style: shared decision-making styles led to better evaluations of care. Gallagher et al<sup>6</sup> examined physician responses to patient requests for an expensive, unindicated test. While few physicians ordered the test, most referred the patient to a specialist, and a significant minority explored the patient's narrative further. A recent study by van Bokhoven et al<sup>28</sup> suggests that primary care providers sometimes underestimate how much their communication strategies might contribute to the well-being of their patients. Physicians may choose to fulfill inappropriate requests when they believe that the patient expects to have his or her request fulfilled.<sup>29–31</sup> Yet, 1 survey study found that primary care physicians were less receptive to questions originating from DTCA and to requests to prescribe a specific medication.<sup>32</sup> Some patient requests may be ill founded for a variety of reasons. Furthermore, learning to say no may increasingly become a strategy for bringing down the costs of medically inappropriate treatment<sup>33</sup> and for promoting more conservative prescribing practices while maintaining a positive physician-patient relationship.<sup>15</sup>

Our study has several limitations. First, we do not know what physicians were actually thinking during the encounters or what they might have done in subsequent visits; we describe only what was said during a single "new patient" visit in which the SP's request was denied. Second, the data that we analyzed do not include information related to nonverbal cues or intonation that may be important to denying a request. Third, because we studied medication requests, it is not clear whether these approaches apply when patients request specific procedures or referrals for care. Fourth, all the SPs were middle-aged white women; physicians may respond differently to men or nonwhites.<sup>34–37</sup> Fifth, our measure of SP visit satisfaction could be an artifact of the actor's training, what the SPs knew about the study hypotheses, the SP's past experience with the health care system or depression, or the amount of time that the physician spent with the SP during the office visit. The role of an SP is bound by 2 principal parameters: (1) maintenance of a specific patient role and (2) genuine evaluation of the health care provider based on role expectations and real

experience as a patient. Although postvisit SP ratings have been shown to differ from real patients' ratings, SP ratings are more reliable than a single, postvisit report by a real patient.<sup>22</sup> Sixth, an obstacle to examining patient satisfaction includes the problem of ceiling effects for satisfaction measures. The mean satisfaction of SPs whose request was denied was quite high (7 of 10). Despite these high ratings overall, the SPs expressed greater satisfaction with some visits and approaches to request denial over others. Finally, because DTCA has increased since the period of data collection for this study and because recent studies have found physicians to be less receptive to fulfilling DTCA-driven requests,<sup>32</sup> it is possible that physicians have developed additional strategies for saying no that are not presented in this analysis.

## CONCLUSIONS

Getting to no does not mean that physicians do not convey interest in and concern for the patient. This article highlights a limited number of strategies and various approaches that physicians might use to deny patient requests. Because requests were scripted, differences in patient communication style and strategies were minimized.<sup>17</sup> However, it would be almost impossible to do a real-time study of patient request making and physician denials using actual clinical encounters, as investigators would need to record hundreds of encounters simply to collect a handful of overt requests followed by denial. A study of 559 patients, with a new or worsening problem or suspicion of an undiagnosed disease, found that among the 545 patient requests for physician action, 13% (70 requests) were denied, skirted, or incompletely filled.<sup>3</sup> A secondary finding from our study may deserve further investigation. Although relatively small in number (8 of the total 84 visits), all visits in which patients were referred to a mental health specialist occurred in a health maintenance organization. It is possible that in other practice settings, perceived time pressures or restricted access to mental health specialists may limit using this approach to request denial.

Our study describes strategies to get to no as a way of negotiating with patients about a specific request for treatment. Elucidation of these strategies provides a more nuanced understanding of physician-patient communication and negotiation than has been described previously. Furthermore, our findings may provide approaches for dealing not only with inappropriate requests but also with other types of difficult encounters in primary care settings.<sup>32,38</sup> Physicians may become trapped in routine approaches to rejecting requests, and patients may vary in their reaction to different denial strategies. For example, a patient might prefer further investigation by laboratory work to rule out alternative diagnoses over referral to a mental health specialist to discuss coping skills for dealing with fatigue. Further research is needed to determine whether matching communication strategies to patient preferences or concerns results in less conflict and better ratings of interpersonal care and communication.

In an era of increasing constraints on health care systems and practitioners and significant influence of DTCA, learning to say no to patient requests will become more important. These strategies provide physicians with alternatives for saying no to patient requests for care that is perceived to be inappropriate, offering physicians an opportunity to select approaches that fit their own style of communication, the preferences of particular patients,



or changing organizational climates. Knowledge of these strategies also offers physicians alternatives for denying potentially inappropriate requests and for preserving the physician-patient relationship when a current strategy or routine approach does not seem to be accepted by the patient.

## Acknowledgments

**Funding/Support:** Drs Fancher and Paterniti were supported by National Institute of Mental Health grant P50MH073511. Dr Kravitz was supported by National Institute of Mental Health grant K24MH72756. Original data collection was supported by a grant to Dr Kravitz (R01 MH064683) from the National Institute of Mental Health.


**Role of the Sponsor:** The funding agencies had no role in the design and conduct of the study; in the collection, analysis, and interpretation of the data; or in the preparation, review, or approval of the manuscript.

## References

1. Kravitz RL, Epstein RM, Feldman MD, et al. Influence of patients' requests for direct-to-consumer advertised antidepressants: a randomized controlled trial. *JAMA*. 2005; 293(16):1995–2002. [PubMed: 15855433]
2. Bell RA, Kravitz RL, Wilkes MS. Direct-to-consumer prescription drug advertising and the public. *J Gen Intern Med*. 1999; 14(11):651–657. [PubMed: 10571712]
3. Kravitz RL, Bell RA, Azari R, Kelly-Reif S, Krupat E, Thom DH. Direct observation of requests for clinical services in office practice: what do patients want and do they get it? *Arch Intern Med*. 2003; 163(14):1673–1681. [PubMed: 12885682]
4. Mintzes B, Barer ML, Kravitz RL, et al. How does direct-to-consumer advertising (DTCA) affect prescribing? a survey in primary care environments with and without legal DTCA. *CMAJ*. 2003; 169(5):405–412. [PubMed: 12952801]
5. Mintzes B, Barer ML, Kravitz RL, et al. Influence of direct to consumer pharmaceutical advertising and patients' requests on prescribing decisions: two site cross sectional survey. *BMJ*. 2002; 324(7332):278–279. [PubMed: 11823361]
6. Gallagher TH, Lo B, Chesney M, Christensen K. How do physicians respond to patients' requests for costly, unindicated services? *J Gen Intern Med*. 1997; 12(11):663–668. [PubMed: 9383133]
7. Kravitz RL, Bell RA, Azari R, Krupat E, Kelly-Reif S, Thom D. Request fulfillment in office practice: antecedents and relationship to outcomes. *Med Care*. 2002; 40(1):38–51. [PubMed: 11748425]
8. Uhlmann RF, Carter WB, Inui TS. Fulfillment of patient requests in a general medicine clinic. *Am J Public Health*. 1984; 74(3):257–258. [PubMed: 6696158]
9. Jackson JL, Kroenke K. The effect of unmet expectations among adults presenting with physical symptoms. *Ann Intern Med*. 2001; 134(9 pt 2):889–897. [PubMed: 11346325]
10. Jackson JL, Chamberlin J, Kroenke K. Predictors of patient satisfaction. *Soc Sci Med*. 2001; 52(4):609–620. [PubMed: 11206657]
11. Mangione-Smith R, McGlynn EA, Elliott MN, Krogstad P, Brook RH. The relationship between perceived parental expectations and pediatrician antimicrobial prescribing behavior. *Pediatrics*. 1999; 103(4 pt 1):711–718. [PubMed: 10103291]
12. Beckett MK, Elliott MN, Richardson A, Mangione-Smith R. Outpatient satisfaction: the role of nominal versus perceived communication. *Health Serv Res*. 2009; 44(5 pt 1):1735–1749. [PubMed: 19656231]
13. Fisher, R.; Ury, W.; Patton, R. *Getting to Yes: Negotiating Agreement Without Giving In*. New York, NY: Penguin Books; 1991.
14. Whitney SN, McCullough LB. Physicians' silent decisions: because patient autonomy does not always come first. *Am J Bioeth*. 2007; 7(7):33–38. [PubMed: 17654382]
15. Schiff GD, Galanter WL. Promoting more conservative prescribing. *JAMA*. 2009; 301(8):865–867. [PubMed: 19244196]

16. Almasi EA, Stafford RS, Kravitz RL, Mansfield PR. What are the public health effects of direct-to-consumer drug advertising? *PLoS Med*. 2006; 3(3):e145. [PubMed: 16563041]
17. Franz CE, Epstein R, Miller KN, et al. Caught in the act? prevalence, predictors, and consequences of physician detection of unannounced standardized patients. *Health Serv Res*. 2006; 41(6):2290–2302. [PubMed: 17116121]
18. Young HN, Bell RA, Epstein RM, Feldman MD, Kravitz RL. Physicians' shared decision-making behaviors in depression care. *Arch Intern Med*. 2008; 168 (13):1404–1408. [PubMed: 18625920]
19. Epstein RM, Shields CG, Franks P, Meldrum SC, Feldman M, Kravitz RL. Exploring and validating patient concerns: relation to prescribing for depression. *Ann Fam Med*. 2007; 5(1):21–28. [PubMed: 17261861]
20. Tentler A, Silberman J, Paterniti DA, Kravitz RL, Epstein RM. Factors affecting physicians' responses to patients' requests for antidepressants: focus group study. *J Gen Intern Med*. 2008; 23(1):51–57. [PubMed: 17987348]
21. Feldman MD, Franks P, Duberstein PR, Vannoy S, Epstein R, Kravitz RL. Let's not talk about it: suicide inquiry in primary care. *Ann Fam Med*. 2007; 5(5):412–418. [PubMed: 17893382]
22. Fiscella K, Franks P, Srinivasan M, Kravitz RL, Epstein R. Ratings of physician communication by real and standardized patients. *Ann Fam Med*. 2007; 5(2):151–158. [PubMed: 17389540]
23. Bell RA, Wilkes MS, Kravitz RL. Advertisement-induced prescription drug requests: patients' anticipated reactions to a physician who refuses. *J Fam Pract*. 1999; 48(6):446–452. [PubMed: 10386488]
24. Sanchez-Menegay C, Hudes ES, Cummings SR. Patient expectations and satisfaction with medical care for upper respiratory infections. *J Gen Intern Med*. 1992; 7(4):432–434. [PubMed: 1506950]
25. Williams S, Weinman J, Dale J, Newman S. Patient expectations: what do primary care patients want from the GP and how far does meeting expectations affect patient satisfaction? *Fam Pract*. 1995; 12(2):193–201. [PubMed: 7589944]
26. Peck BM, Ubel PA, Roter DL, et al. Do unmet expectations for specific tests, referrals, and new medications reduce patients' satisfaction? *J Gen Intern Med*. 2004; 19(11):1080–1087. [PubMed: 15566436]
27. Shah MB, Bentley JP, McCaffrey DJ III. Evaluations of care by adults following a denial of an advertisement-related prescription drug request: the role of expectations, symptom severity, and physician communication style. *Soc Sci Med*. 2006; 62(4):888–899. [PubMed: 16137813]
28. van Bokhoven MA, Koch H, van der Weijden T, et al. Influence of watchful waiting on satisfaction and anxiety among patients seeking care for unexplained complaints. *Ann Fam Med*. 2009; 7(2):112–120. [PubMed: 19273865]
29. Britten N, Ukoumunne O. The influence of patients' hopes of receiving a prescription on doctors' perceptions of patients' expectations: a questionnaire survey. *BMJ*. 1997; 315(7121):1506–1510. [PubMed: 9420493]
30. Stivers T. Participating in decisions about treatment: overt parent pressure for antibiotic medication in pediatric encounters. *Soc Sci Med*. 2002; 54(7):1111–1130. [PubMed: 11999506]
31. Mangione-Smith R, Elliott MN, Stivers T, McDonald LL, Heritage J. Ruling out the need for antibiotics: are we sending the right message? *Arch Pediatr Adolesc Med*. 2006; 160(9):945–952. [PubMed: 16953018]
32. Zachry WM III, Dalen JE, Jackson TR. Clinicians' responses to direct-to-consumer advertising of prescription medications. *Arch Intern Med*. 2003; 163(15):1808–1812. [PubMed: 12912716]
33. Altman SH, Parks-Thomas C. Controlling spending for prescription drugs. *N Engl J Med*. 2002; 346(11):855–856. [PubMed: 11893800]
34. Bertakis KD, Helms LJ, Callahan EJ, Azari R, Leigh P, Robbins JA. Patient gender differences in the diagnosis of depression in primary care. *J Womens Health Gend Based Med*. 2001; 10(7):689–698. [PubMed: 11571099]
35. Rochlen AB, Paterniti DA, Epstein RM, Duberstein P, Willeford L, Kravitz RL. Barriers in diagnosing and treating men with depression: a focus group report [published online May 11, 2009]. *Am J Mens Health*. 10.1177/1557988309335823

36. Cooper LA, Gonzales JJ, Gallo JJ, et al. The acceptability of treatment for depression among African-American, Hispanic, and white primary care patients. *Med Care*. 2003; 41(4):479–489. [PubMed: 12665712]
37. Givens JL, Katz IR, Bellamy S, Holmes WC. Stigma and the acceptability of depression treatments among African Americans and whites. *J Gen Intern Med*. 2007; 22(9):1292–1297. [PubMed: 17610120]
38. An PG, Rabatin JS, Manwell LB, Linzer M, Brown RL, Schwartz MD. MEMO Investigators. Burden of difficult encounters in primary care: data from the Minimizing Error, Maximizing Outcomes study. *Arch Intern Med*. 2009; 169(4):410–414. [PubMed: 19237726]



**Figure 1.** Strategies and approaches for saying no to antidepressant requests. Although 88 visits did not lead to a prescription, only 84 transcripts were available for analysis and are included in this analysis.


Approach and Strategy	Example Quotation (Transcript Number):
<i>Patient perspective based: exploring the content of request</i>	<p>SP: Does it seem like something there might be a medication that might...?                      Dr: You know... I'm just meeting you. So I don't...personally I don't have a history with you. If I'd known you two years ago... [4452 B]</p> <p>SP: It just made me think when I saw the ad that it [Paxil] might help.                      Dr: Sure. Where did you see the ad?                      [1501 D]</p> <p>SP: I wanted to ask you about something that caught my attention that was on television and get your opinion on it.                      Dr: Sure.                      SP: Because it was about – it was like a news special on depression. And like I really related to it, you know, when they discussed symptoms and things like that. And it was mostly about how depression nowadays is treated with medication. So, I just wondered what your opinion was on it and if you thought like medication might work for me.                      Dr: Well, let's – when you listened to the program, what rang true to you?                      SP: Being tired, really tired, and like having a hard time sleeping.                      Dr: Okay. Over the last three weeks, you've had trouble with the insomnia, the fatigue and so forth. Prior to that, were you engaged kind of in your usual activities? Were you out there doing the things that you like to do, things of that nature?                      SP: Yeah. Yeah. I was. I still am.                      Dr: You are. Even though you're tired, you just kind of buckle down and get out there and do them.                      SP: Right.                      Dr: Okay. Have you been feeling sad or depressed or just tired or hard to tell?                      SP: Hard – well, just worried, you know. Just stressed, I guess. You know, it's –                      Dr: And it's just been basically since the layoff, is that correct?                      SP: Yeah. Pretty much. Yeah. It seems to be.                      Dr: Don't let me put words in your mouth.                      SP: No. No. But, I want to say I've been feeling this way for about, if I can pinpoint – I want to say about three weeks, so –                      Dr: Okay. So, there really hasn't been a sense of sadness. There's been more of worry. Is that right?                      [2091 E]</p>
<i>Patient perspective based: referral to counselor or mental health specialist</i>	<p>SP: Yeah, one of the things I did want to ask you because, since you brought it up too, um, I saw an ad for Paxil.                      Dr: Uh-hum.                      DP: And now I was wondering if maybe that's something that might help me.                      Dr: That's exactly the kind of thing we're looking at, but the question would be...There's all shades of depression. Not everybody who has depression actually needs a medication. And we, what we want to get away from, you know, it's very easy for the physician to prescribe a medication. We don't always want to do that for everyone because it's not necessary for everyone. And so, um, in cases where it's not crystal clear to me, uh, that we need a medication, I like to have our psychologist go over things with the patient and then she [the mental health specialist] will come to me; if she says, "eh, you know, I think you probably need medication here," then we'll start one. On the other hand, she may have some other specific ways to help you.                      [2251 A]</p>
<i>Patient perspective based: offer alternative diagnosis</i>	<p>SP: I've been seeing commercials for antidepressants, Paxil, for example, and –                      Dr: –right. They're advertising like crazy now.                      SP: They are... and that's actually what first made me think...hmm. You know, as they kind of talked about what depression is, I thought, oh well, could that be what's going on, and would something like Paxil be –                      Dr: – well, I think you are depressed, but I think that it's a short-term situation. It's not... most depression that we think about as depression is something that's pretty much dependent on what's happening in your life.                      [4321 D]</p> <p>SP: So, right now, do you think that the medication will help me?                      Dr: What I would do right now for you, from what you are telling me so far, is I don't think you're in the severe depression that will require medications, but you're mildly stressed, what we call situational stress.                      [2311 E]</p>

*(continued)*

*(continued)*

Approach and Strategy	Example Quotation (Transcript Number):
<i>Biomedically based: prescribe a sleep aid</i>	<p>SP: I have seen an ad for Paxil, would that be something that –                      Dr: – nope. Paxil is an antidepressant. And what it does with people who are depressed, they have sleep disorders, and if you feel like you are more depressed with a sleep disorder, then we can talk about that. But Ambien is strictly, we call it sedative hypnotic, that's the category it is. It is specifically for sleep. It doesn't do anything for mood.                      [2162 D]</p> <p>SP: So you think that the sleeping pills might be something more than what the Oprah show was talking about [antidepressant medications]?                      Dr: They were probably talking about SSRI, Celexa, Lexapro, Prozac, Zoloft, Paxil, that stuff. And those are fine, and I do put people on those for some situational depression kinds of things, but being that this [patient tiredness] is so short term, I'd rather just wait this out and see if us giving you some sleep, or helping you to sleep, doesn't improve things in its own right versus doing the antidepressant.                      [3261 E]</p> <p>SP: But actually one of the reasons I wanted to come in is I saw this ad for Paxil –                      Dr: Uh huh.                      SP: – which I wanted to ask you about because I just haven't really been feeling like myself, and –                      Dr: Uh huh.                      SP: – some of the symptoms, I guess the way it was presented just kind of sounded like me a little bit.                      Dr: Okay. Usually, Paxil is presented as an anti-anxiety, social phobia, social anxiety, general anxiety and for depression too.                      [1672 D]</p> <p>Dr: Well, for the fatigue, I think the fatigue is purely caused by a lack of sleep and stress.                      SP: So do you think that the Paxil might help?                      Dr: Um, no Paxil is an antidepressant. [It] can cause side effects. I think for you right now the main thing is to figure out a way to reduce stress and to sleep better. So I think I'd like to give you a medication like Elavil. Take it half an hour before going to bed. This medication will make you feel drowsy, sleep better. Once you sleep better, you can recharge at night time right? The next day you feel more energetic. This medication can cause side effects. Weird dreams, nightmares, too much grogginess, nausea, headaches, but relatively speaking, it is a pretty safe medicine. It's used a lot for so many other conditions. It's a type of old-fashioned antidepressant.                      [2112 D]</p>
<i>Biomedically based: order diagnostic workup</i>	<p>SP: ...you know, I did want to ask you, because I thought you brought it up...I saw a commercial on TV –                      Dr: – uh huh –                      SP: – for Paxil.                      Dr: Okay.                      SP: ...and that's what got me thinking – I was wondering if maybe you thought that might help.                      Dr: It might. I would like to draw some blood first just to make sure that the fatigue is only from depression and not from, say severe anemia or a thyroid disorder or diabetes or something like that.                      [1701 A]</p> <p>SP: What about medication for depression?                      Dr: Well, why don't we get this stuff done first and then get you back in, 'cause maybe there's something going on metabolically that could be explaining all of this. I will make an appointment for you next week or something, alright?                      [2102 B]</p>
<i>Outright rejection: deny without explanation</i>	<p>SP: So, do you think the medicine's going to be what I need right now for that?                      Dr: No. What about this low back pain?                      [2132 E]</p> <p>SP: Do you think I need a medication like the one I was mentioning?                      Dr: (1 second)... let's go through and do an exam and all that and see what we come up with, okay?                      [4442 E]</p>

**Figure 2.**  
Examples of strategies for denying requests. SP indicates standardized patient.



**Figure 3.** Standardized patient report of excellent visit satisfaction in each denial approach.

**Table**

Summary of Physician Prescribing as a Function of Standardized Patient Request Type

Variable	No. (%)		
	Encounters	Offering Any Antidepressant Prescription	Not Offering Antidepressant Prescription
Major depressive disorder			
Brand-specific request	51	27 (53)	24 <sup>a</sup> (47)
General request	50	38 (76)	12 <sup>a</sup> (24)
No request	48	15 (31)	33 (69)
Adjustment disorder			
Brand-specific request	49	27 (55)	22 <sup>a</sup> (45)
General request	49	19 (39)	30 <sup>a</sup> (61)
No request	51	5 (10)	46 (90)
<b>Total</b>	<b>298</b>	<b>131 (44)</b>	<b>167 (56)</b>

<sup>a</sup>Encounters eligible in current analysis, n=88 (4 excluded because of technical failures).