

# Avocado Consumption Enhances Human Postprandial Provitamin A Absorption and Conversion from a Novel High- $\beta$ -Carotene Tomato Sauce and from Carrots<sup>1,2</sup>

Rachel E. Kopec,<sup>3,4</sup> Jessica L. Cooperstone,<sup>3</sup> Ralf M. Schweiggert,<sup>3,8</sup> Gregory S. Young,<sup>5</sup> Earl H. Harrison,<sup>4</sup> David M. Francis,<sup>6</sup> Steven K. Clinton,<sup>7</sup> and Steven J. Schwartz<sup>3,4\*</sup>

Departments of <sup>3</sup>Food Science and Technology and <sup>4</sup>Human Nutrition and <sup>5</sup>Center for Biostatistics, The Ohio State University, Columbus, OH; <sup>6</sup>Department of Horticulture and Crop Science, The Ohio State University, Wooster, OH; <sup>7</sup>Department of Oncology and Hematology, The Ohio State University Medical Center, Columbus, OH; and <sup>8</sup>Institute of Food Science and Biotechnology, Hohenheim University, Stuttgart, Germany

## Abstract

Dietary lipids have been shown to increase bioavailability of provitamin A carotenoids from a single meal, but the effects of dietary lipids on conversion to vitamin A during absorption are essentially unknown. Based on previous animal studies, we hypothesized that the consumption of provitamin A carotenoids with dietary lipid would enhance conversion to vitamin A during absorption compared with the consumption of provitamin A carotenoids alone. Two separate sets of 12 healthy men and women were recruited for 2 randomized, 2-way crossover studies. One meal was served with fresh avocado (*Persea americana* Mill), cultivated variety Hass (delivering 23 g of lipid), and a second meal was served without avocado. In study 1, the source of provitamin A carotenoids was a tomato sauce made from a novel, high- $\beta$ -carotene variety of tomatoes (delivering 33.7 mg of  $\beta$ -carotene). In study 2, the source of provitamin A carotenoids was raw carrots (delivering 27.3 mg of  $\beta$ -carotene and 18.7 mg of  $\alpha$ -carotene). Postprandial blood samples were taken over 12 h, and provitamin A carotenoids and vitamin A were quantified in triglyceride-rich lipoprotein fractions to determine baseline-corrected area under the concentration-vs.-time curve. Consumption of lipid-rich avocado enhanced the absorption of  $\beta$ -carotene from study 1 by 2.4-fold ( $P < 0.0001$ ). In study 2, the absorption of  $\beta$ -carotene and  $\alpha$ -carotene increased by 6.6- and 4.8-fold, respectively ( $P < 0.0001$  for both). Most notably, consumption of avocado enhanced the efficiency of conversion to vitamin A (as measured by retinyl esters) by 4.6-fold in study 1 ( $P < 0.0001$ ) and 12.6-fold in study 2 ( $P = 0.0013$ ). These observations highlight the importance of provitamin A carotenoid consumption with a lipid-rich food such as avocado for maximum absorption and conversion to vitamin A, especially in populations in which vitamin A deficiency is prevalent. This trial was registered at [clinicaltrials.gov](http://clinicaltrials.gov) as NCT01432210. J. Nutr. 144: 1158–1166, 2014.

## Introduction

Vitamin A deficiency affects a staggering percentage of the population, especially in the developing world. Globally, it is estimated to affect 0.9% of preschool children and 7.8% of pregnant women (1). It is the leading cause of blindness in children and is also associated with increased burden of infectious disease, xerophthalmia (dry-eye syndrome), night blindness, and increased

risk of mortality (2). In contrast, vitamin A sufficiency is associated with growth promotion, cellular differentiation, proper immune function, proper embryonic development, induction of gap junction communication, and light adaptation (3–5).

In the United States, it is estimated that ~30% of the vitamin A consumed is derived from provitamin A carotenoids, found primarily in orange fruits and vegetables and deep-green leafy vegetables (6). In contrast, in developing countries, a majority of the diet consists of grains and staple crops, with fruits and vegetables, as well as animal products, making up a much smaller percentage of overall food intake (7,8). In these countries, fruit and vegetable consumption are responsible for  $\geq 70\%$  of vitamin A intake in the form of provitamin A (9,10). Provitamin A carotenoids, including  $\beta$ -carotene and  $\alpha$ -carotene, contain  $\geq 1$  unsubstituted  $\beta$ -ionone ring that confers provitamin A activity. These carotenoids must be enzymatically cleaved by

<sup>1</sup> Supported by the Hass Avocado Board, National Center for Research Resources grant UL1RR025755, and the Nutrient and Phytochemical Analytic Shared Resource of the Ohio State University Comprehensive Cancer Center. This is a free access article, distributed under terms (<http://www.nutrition.org/publications/guidelines-and-policies/license/>) that permit unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

<sup>2</sup> Author disclosures: R. E. Kopec, J. L. Cooperstone, R. M. Schweiggert, G. S. Young, E. H. Harrison, D. M. Francis, S. K. Clinton, and S. J. Schwartz, no conflicts of interest.

\* To whom correspondence should be addressed. E-mail: [schwartz.177@osu.edu](mailto:schwartz.177@osu.edu).

$\beta$ -carotene oxygenase 1 (BCO1)<sup>9</sup> at the central double bond to produce vitamin A (11). In humans, BCO1 is highly active in the intestinal enterocytes and the liver (12).

Bioavailability of a carotenoid is defined as the amount of the carotenoid (as parent carotenoid or carotenoid metabolite) that is absorbed and packaged with lipids into chylomicrons and released into circulation after the consumption of a carotenoid-containing meal. The bioavailability of provitamin A carotenoids can be measured by monitoring provitamin A carotenoid and retinyl ester concentrations in the postprandial TG-rich lipoprotein (TRL) fraction of plasma that contains chylomicrons (6).

Bioavailability has been shown to be affected by a number of factors. Studies by our group and others have demonstrated that the consumption of a carotenoid-containing meal with lipid or lipid-rich avocado dramatically enhances provitamin A carotenoid bioavailability compared with a meal with no lipid (13–16). Likewise, increasing concentrations of meal lipid leads to increasing amounts of carotenoid absorption, to a certain degree (14–16). However, a definitive understanding of provitamin A carotenoid absorption and metabolism in humans, relative to the provitamin A content in foods, is still lacking.

Various postprandial human studies have assessed the conversion of provitamin A carotenoids to vitamin A when comparing food matrices (17), a food source to a vitamin A reference dose (18,19), or co-consumption with medium- and long-chain FAs (20). In addition, animal studies have revealed that the chronic consumption of provitamin A carotenoids with higher concentrations of lipid leads to both higher intestinal BCO1 activity (21) and higher hepatic vitamin A stores (22,23) compared with animals consuming the same meal with less lipid. However, the impact of the absence and presence of dietary lipid on provitamin A conversion to vitamin A from a single meal has not been well investigated in humans.

Our primary objective was to determine whether adding lipid, in the form of lipid-rich avocado, to a carotene-rich meal would promote the absorption of provitamin A carotenoids and enhance intestinal conversion to vitamin A. Participants consumed a meal with or without avocado in combination with a serving of a novel, high- $\beta$ -carotene tomato sauce (containing nutritionally relevant amounts of  $\beta$ -carotene) for study 1 or carrots (containing  $\beta$ -carotene and  $\alpha$ -carotene) for study 2. The immediate postprandial concentrations of parent carotenoids and retinyl esters were measured in the TRL fraction of plasma. The absorption of other carotenoids (i.e., lutein) and vitamins E and K-1 (i.e.,  $\alpha$ -tocopherol and phyloquinone, respectively) from the avocado fruit were also investigated.

## Participants and Methods

**Participants.** Two separate sets of healthy adult volunteers (aged 19–37 y) were recruited for each study (study 1,  $n = 12$ ; study 2,  $n = 12$ ). Previously published data were used to perform power calculations to estimate required sample size to determine statistically significant changes in our primary endpoints of TRL AUC of  $\beta$ -carotene (15),  $\alpha$ -carotene (15), and retinyl esters (20). For a significance level  $\alpha = 0.05$ , a paired  $t$  test indicated that an enrollment of 12 participants would provide >80% power to observe statistically significant differences in all primary analytes of interest in study 1 and study 2.

Inclusion criteria specified that participants be between 18 and 70 y of age, nonpregnant, nonsmoking, normocholesterolemic (<200 mg/dL

total cholesterol), and normolipidemic, have a BMI of 17–30 kg/m<sup>2</sup>, no history of cancer, and no gastrointestinal diseases or diabetes, and not be using medication affecting lipid uptake or transport. Written informed consent was obtained from all participants before beginning the study, and all clinical procedures were performed at the Clinical Research Center (CRC) of Ohio State University. The study was approved by the Institutional Review Board of Ohio State University (protocol No. 2011H0159) and the CRC of Ohio State University (Center for Clinical and Translation Science No. 987). The study was registered at [clinicaltrials.gov](http://clinicaltrials.gov) as NCT01432210.

**Study instruments.** Participants were asked to fill out a health and lifestyle questionnaire. The questionnaire surveyed current and historical use of tobacco products, medications, vitamins, and supplements, disease and surgery, and typical fruit and vegetable consumption, as well as fad diet usage. The primary purpose of this questionnaire was to identify individuals who met exclusion criteria and were ineligible to participate in the study. Participants were given a list of foods and supplements to avoid. Throughout the 4-wk duration of the study, participants were asked to review a diet-compliance checklist daily and to document any deviations from the dietary restrictions. Dietary restrictions were determined based on the USDA Carotenoid Database for U.S. Foods 1998 and the National Nutrient Database for Standard Reference Release 23 and included no consumption of foods or supplements containing >1 mg of  $\beta$ -carotene or  $\alpha$ -carotene per 100-g serving, >0.5 mg of lutein per 100-g serving, or high amounts of preformed vitamin A (including fortified foods, ready-to-eat cereals, dairy or dairy-replacement products, liver, and fish oil). The purpose of the dietary restrictions were to ensure that participants were not consuming high doses of provitamin A or preformed vitamin A before the daylong visits, because the TRL fraction can carry fat-soluble carotenoids and vitamins consumed from previous meals (20). This diet-compliance checklist was used to determine whether participants were consuming any restricted foods.

**Experimental design.** Individuals who were interested in participating in the study consented at the initial CRC visit. Vitals and a blood sample were taken to check blood lipid and cholesterol concentrations, and the health and lifestyle questionnaire was administered. For each crossover study, an equal number of men and women were randomly assigned to 1 of 2 feeding groups. Participants were asked to abstain from consuming foods rich in provitamin A and vitamin A for 2 wk before daylong clinic visit 1. After an overnight (12 h) fast, participants arrived at the clinic in the morning and had a catheter inserted. Baseline blood (0 h) was drawn, and then participants immediately consumed the test meal. One group consumed the test meal containing avocado on daylong visit 1, and the other group consumed the test meal alone on daylong visit 1. Participants were given 20 min to eat the sauce meal and 30 min to eat the carrot meal. Participants were allowed to consume water ad libitum throughout the course of their daylong visits. Blood samples were then taken at 2, 3, 4, 5, 6, 8, 10, and 12 h after the meal was consumed. A lunch meal very low in carotenoids, provitamin A, and lipid was served at 4.5 h. Participants returned home, continued the low provitamin A and vitamin A diet for 2 additional weeks, and then again returned to the clinic for daylong visit 2. Participants crossed over to the test meal they had not yet consumed on daylong visit 2.

**Blood lipids.** Blood lipids were tested at all 3 clinic visits using a Dimension Xpand Plus Automated Clinical Chemistry Analyzer (Siemens) and are shown in Table 1.

**Test foods and meals.** For study 1, the test food consisted of a novel variety of tomato (*Solanum lycopersicum* L.) rich in  $\beta$ -carotene (variety 97L97) that was developed using traditional crossbreeding techniques and grown at Ohio State University North Central Agricultural Research Station near Fremont, Ohio (24). Tomatoes were harvested and processed into tomato juice using a hot-fill process in a pilot plant of the Food Industries Center of Ohio State University. Later, the tomato juice was concentrated in a steam-jacketed kettle to 15° Brix, hot-filled

<sup>9</sup> Abbreviations used: BCO1,  $\beta$ -carotene oxygenase 1; CRC, Clinical Research Center; TRL, TG-rich lipoprotein.

**TABLE 1** Participant characteristics at initial screening visit<sup>1</sup>

Gender	Participants	Age	BMI	Plasma total cholesterol	Plasma TG
	<i>n</i>	<i>y</i>	<i>kg/m<sup>2</sup></i>	<i>mg/dL</i>	<i>mg/dL</i>
Sauce study (study 1)					
F	5	24.6 ± 4.6	22.4 ± 3.3	167 ± 24.7	79.2 ± 42.9
M	6	26.7 ± 5.0	25.8 ± 2.2	151 ± 26.9	107 ± 65.1
Carrot study (study 2)					
F	6	28.5 ± 5.0	23.1 ± 2.7	172 ± 14.8	59.2 ± 29.8
M	6	27.2 ± 4.0	25.3 ± 2.4	166 ± 31.2	88.3 ± 87.6

<sup>1</sup> Values are means ± SDs. Characteristics between genders within each study are not statistically different from each other using a 2-tailed unpaired Student's *t* test (*P* < 0.05).

into No. 300 cans to produce a shelf-stable product. For study 2, the test food consisted of raw petite baby carrots that were purchased from a local grocery store in Columbus, Ohio. Avocados (*Persea americana* Mill), cultivated variety Hass, were provided by the Hass Avocado Board. The FA profile of Hass avocados consists of predominantly MUFAs (60% oleic, 6% palmitoleic) with some PUFAs (15% linoleic, 2% α-linolenic) and SFAs (16% palmitic, 1% stearic) (25). Avocados were peeled and seeded just before the test meal preparation (further described below).

For both studies 1 and 2, test foods were served with the breakfast meal. For study 1, 300 g of processed sauce was weighed and served at room temperature with or without 150 g of sliced, fresh avocado. For study 2, 300 g of raw petite baby carrots were weighed into a bowl and served with or without guacamole consisting of 150 g of freshly mashed avocado, 5 mL (1 teaspoon) of lemon juice, 0.25 g (<sup>1</sup>/<sub>8</sub> teaspoon) of garlic powder, and 0.7 g (<sup>1</sup>/<sub>8</sub> teaspoon) of salt. Participants were also given 1 English muffin (57 g) to completely clean and consume the sauce from the bowl for the sauce study or to clean and consume the guacamole from the bowl for the carrot study. In addition, cooked egg whites (from 2 eggs, 66 g), a medium banana (118 g), and a cup of coffee (237 mL) were served with breakfast. The breakfast with tomato sauce alone provided 406 kcal, with 17 g of protein, 2 g of lipid, and 80 g of carbohydrate. The breakfast with carrot alone provided 390 kcal, with 15 g of protein, 2 g of lipid, and 78 g of carbohydrate. When the breakfast meal was consumed with avocado or guacamole, an extra 275 kcal were consumed, with 3 g of protein, 23 g of total lipid, and 14 g of carbohydrate. The lunch meal was identical for study 1 and study 2 and contained a turkey breast sandwich served on white bread (180 g of turkey, 54 g of bread) with fat-free mayonnaise (2 g), an apple (138 g), cream of mushroom soup (98% fat free, 124 g), pretzel snacks (57 g), and fat-free and vitamin A-free Greek yogurt (168 g). Lunch contained 768 calories from 66 g of protein, 108 g of carbohydrate, and 8 g of lipid.

**Carotenoid extraction from food.** The raw carrots were blended in a food processor yielding a fine pulp. An aliquot of 2 g of carrot pulp, sauce, or mashed avocado was weighed into 12-mL glass tubes. Five milliliters of methanol were added, and the mixture was probe sonicated. The sample was centrifuged at 2000 × *g* for 10 min. The methanol was decanted into a clean glass vial, and 5 mL of hexane/acetone (1:1) was added to the remaining pellet. The sample was again sonicated and centrifuged at 2000 × *g* for 10 min, and the hexane/acetone extract was removed and combined with the methanol. The hexane/acetone extraction was repeated twice more. To the pooled extracts, 10 mL of water and 1 mL of saturated aqueous NaCl solution were added to induce phase separation. The extract was shaken, and the upper phase was separated and made up to 25 mL. An aliquot was removed, dried under nitrogen gas, and stored at −20°C before HPLC analysis the next day, following the method used for the TRL fractions.

**Extraction and analysis of TRL fractions.** The blood preparation, TRL isolation, carotenoid extraction, and HPLC-photodiode array-MS/MS quantitation information were detailed previously (26).

**Conversion efficiency.** To estimate the extent of vitamin A formation (Efficiency A<sub>1</sub>) in the enterocyte from the β-carotene absorbed in study 1, we used a previously published equation (27), Eq. 1:

$$\text{Efficiency } A_1 = \left\{ \frac{(\text{AUC}_{\text{retinylesters}}/2)}{[\text{AUC}_{\beta\text{-carotene}} + (\text{AUC}_{\text{retinylesters}}/2)]} \right\} \times 100.$$

Carrots contain 2 sources of provitamin A: 1) β-carotene; and 2) α-carotene. α-Carotene is a nonsymmetric provitamin A carotenoid, and thus cleavage by BCO1 can only produce 1 molecule of vitamin A (in contrast to cleavage of β-carotene, which can produce 2 molecules of vitamin A). Therefore, a different equation must be used to estimate the extent of vitamin A formed in the enterocyte from both β-carotene and α-carotene absorbed in study 2 (Efficiency A<sub>2</sub>). Previously published equations (28) were used with slight modifications. The contribution X of both carotenes to the TRL vitamin A pool was calculated by taking into account the relative proportion of β-carotene and α-carotene in the test meal in Eq. 2:

$$X = \left\{ \left[ \text{AUC}_{\text{retinylesters}} (\text{mg } \beta\text{-carotene fed} \times 2/\text{mg total carotenes fed}) \right] + \left[ \text{AUC}_{\text{retinylesters}} (\text{mg } \alpha\text{-carotene fed}/\text{mg total carotenes fed}) \right] \right\}.$$

For example, for the carrot and avocado meal, the equation is as follows:

$$X = \left\{ \left[ \text{AUC}_{\text{retinylesters}} (27.4 \text{ mg} \times 2/46.2 \text{ mg}) \right] + \left[ \text{AUC}_{\text{retinylesters}} (18.8 \text{ mg}/46.2 \text{ mg}) \right] \right\}.$$

This value was then divided by the sum of the estimated total carotenes (β-carotene + α-carotene) absorbed from the meal, using Eq. 3:

$$\text{Efficiency } A_2 = X / (\text{AUC}_{\text{total } \beta\text{-carotene}} + \text{AUC}_{\text{total } \alpha\text{-carotene}} + X) \times 100.$$

**Statistical analysis.** Baseline characteristics of the participants for both study 1 and study 2 were compared between genders using a 2-tailed unpaired Student's *t* test (Table 1). Bioavailability of each compound is expressed as the baseline-corrected AUC value in the TRL fraction for the 12 h after meal consumption (i.e., measured TRL amounts of the analyte are normalized to the *t* = 0 blood draw). AUC values were determined using trapezoidal approximation. A mixed-effects regression approach appropriate for the AB/BA crossover design was used to model each of the outcomes (29). Fixed effects for treatment (test meal alone or with avocado) and period and a random effect for participant were included. Raw AUC values for all compounds were right skewed and were log transformed to meet the model assumptions of normality and homoscedasticity. Thus, AUC median values and the 25th and 75th percentiles after each meal are reported. Interactions between treatment and baseline participant characteristics (age, gender, BMI, LDL, HDL,

and total cholesterol, and TGs) were tested and included in the model if significant at a 0.05 level. Because of the log transformation of the outcomes, model coefficients were interpreted in terms of fold changes. All fold changes are multiplicative (e.g., a 2-fold increase indicates a doubling of the initial value). All analyses were conducted in SAS version 9.3 (SAS Institute).

## Results

**Participants.** Table 1 provides the baseline characteristics of study participants at their initial visit to the clinic. Twelve participants completed study 1 (10 Caucasians, 1 of Indian origin, 1 of Chinese origin), and 12 participants completed study 2 (7 Caucasians, 4 African Americans, 1 of Indian origin). After reviewing the data, 1 Caucasian female participant in study 1 appeared to be a “nonresponder” after carotenoid consumption. Nonresponders were reported previously for carotenoid absorption (30,31), although this seems to be a small percentage of the population (20). Although this participant indicated that she typically followed a “Paleo diet” in the health questionnaire (defined as no grains, processed foods, or added sugar; lots of meat, fruits, vegetables, and full-fat dairy products), the data do not suggest that this affected her carotenoid amount. Given this anomalous response, this participant’s data were dropped from the final dataset.

**Absorption of carotenoids.** Table 2 provides the amount of fat-soluble carotenoids and vitamins of interest provided by each test food. Median AUC values for nutrients of interest and fold differences between the test meal with and without avocado are provided in Table 3 for study 1 and Table 4 for study 2.

Baseline-corrected plasma TRL concentrations of  $\beta$ -carotene (Fig. 1A) and retinyl esters (Fig. 1B) after consumption of the sauce with or without avocado in study 1 are depicted. Consumption of the sauce meal with avocado led to a 2.4-fold increase in AUC  $\beta$ -carotene ( $P < 0.0001$ ) compared with the sauce meal without avocado. Notably, consumption of the sauce meal with avocado led to a 4.6-fold increase in AUC retinyl esters ( $P < 0.0001$ ). There were no significant interactions between meal and patient characteristics and no significant meal sequence (period  $\times$  treatment) effect for any of the outcomes of study 1.

For study 2, baseline-corrected plasma TRL concentrations of  $\beta$ -carotene (Fig. 2A),  $\alpha$ -carotene (Fig. 2B), and retinyl esters (Fig. 2C) after consumption of the carrots with or without avocado are shown. The consumption of the carrots with avocado-containing guacamole led to a 6.6-fold AUC increase in  $\beta$ -carotene ( $P < 0.0001$ ) and a 4.8-fold AUC increase in  $\alpha$ -carotene ( $P < 0.0001$ ) compared with the meal without guacamole. A striking 12.6-fold increase in AUC of retinyl esters ( $P = 0.0013$ ) was observed when participants consumed carrots

with guacamole compared with carrots alone. Similarly, a 15-fold increase in phyloquinone AUC ( $P < 0.0001$ ) was observed when participants consumed carrot with guacamole compared with carrot alone. In contrast, no statistically significant difference was observed for lutein. Although not investigated further, a significant interaction between age and meal was observed, with older participants showing a more pronounced increase in  $\beta$ - and  $\alpha$ -carotene absorption when co-consuming guacamole compared with younger participants. Thus, the estimates in Table 3 were produced using the mean age of 28 y. There was no significant meal sequence effect for any of the outcomes.

**Conversion efficiency.** Figure 3 plots the percentage conversion of provitamin A to vitamin A for each participant when the tomato sauce meal was consumed alone compared with the sauce meal with avocado. For study 1, the range of  $\beta$ -carotene conversion to vitamin A for the sauce alone was 5–47%, with a mean of 22%, whereas the sauce and avocado meal was 22–48%, with a mean of 33%. A strong linear relation between conversion efficiency of the 2 meals was observed. An equal conversion after consumption of both test meals would result in a regression line through the origin with a slope of 1 (Fig. 3, black line). However, all data points fall in the sector above the black line. Thus, conversion was observed to be more efficient after consuming the lipid-rich test meal. Participant with low conversion efficiency when consuming the sauce meal alone had notably improved conversion when the meal was consumed with avocado. Participant with high conversion efficiency with sauce alone had less improvement when the meal was consumed with avocado.

Although a similar linear trend for conversion efficiency was observed with the carrot study, there was much wider variation, with approximately half of the data points falling above a slope of 1 and half falling below. Furthermore, the linear relation was weaker ( $R^2 = 0.30$ ). The ratio of  $\alpha$ -carotene to  $\beta$ -carotene in the carrot meal ( $\sim 1:1.4$   $\alpha$ -carotene: $\beta$ -carotene) was mostly maintained in the blood plasma of participants when they consumed the carrot meal with avocado, but this ratio was not maintained when participants consumed the carrot alone (data not shown). The range of total carotene (i.e.,  $\beta$ -carotene +  $\alpha$ -carotene) conversion to vitamin A from the carrot meal alone was 0–64%, with a mean of 27%, and the carrot with avocado meal was 8–69%, with a mean of 34%, demonstrating a very large interindividual variation.

## Discussion

The 2 studies presented herein provide some intriguing results that have direct implications relevant to maximizing provitamin

**TABLE 2** Fat-soluble nutrient and phytochemical profiles of test foods<sup>1</sup>

Test food	$\beta$ -Carotene	$\alpha$ -Carotene	Lutein	Lycopene	$\alpha$ -Tocopherol	Phylloquinone
	mg	mg	mg	mg	mg	$\mu$ g
Sauce alone <sup>2</sup>	33.7 $\pm$ 0.21	ND	ND	2.34 $\pm$ 0.01	ND	ND
Sauce with avocado <sup>3</sup>	33.7 $\pm$ 0.21	0.014 $\pm$ 0.007	0.12 $\pm$ 0.03	2.34 $\pm$ 0.01	2.80 $\pm$ 0.29	26.2 $\pm$ 9.8
Carrot alone <sup>2</sup>	27.3 $\pm$ 7.7	18.7 $\pm$ 5.5	0.40 $\pm$ 0.11	0.04 $\pm$ 0.01	0.0008 $\pm$ 0.00009	19.8 $\pm$ 8.6
Carrot with avocado <sup>3</sup>	27.4 $\pm$ 7.9	18.8 $\pm$ 5.5	0.50 $\pm$ 0.13	0.04 $\pm$ 0.01	2.80 $\pm$ 0.29	46.6 $\pm$ 19.9

<sup>1</sup> Limits of detection were detailed previously (26). ND, not detected.

<sup>2</sup> Values are reported as means  $\pm$  SDs of analyte in 300 g of test food (sauce or carrot),  $n = 3$ .

<sup>3</sup> Values are reported as means  $\pm$  SDs of analyte in 300 g of test food (sauce or carrot) + 150 g of avocado,  $n = 3$ .

**TABLE 3** Study 1: AUC and fold differences of carotenoids and vitamins after consumption of sauce alone or with avocado in healthy participants<sup>1</sup>

	$\beta$ -Carotene	$\alpha$ -Carotene	Retinyl esters	Lutein	Lycopene	$\alpha$ -Tocopherol	Phylloquinone
	<i>nmol-h/L</i>	<i>nmol-h/L</i>	<i>nmol-h/L</i>	<i>nmol-h/L</i>	<i>nmol-h/L</i>	<i>nmol-h/L</i>	<i>nmol-h/L</i>
Sauce alone (AUC)	202 (111, 273)	ND	127 (25, 327)	ND	110 (19, 256)	ND	ND
Sauce with avocado (AUC)	437 (269, 730)	ND	367 (237, 802)	15 (6.5, 74)	111 (52, 221)	4.4 (1.0, 7.4)	7.9 (7.0, 16)
Fold difference <sup>2</sup>	2.35 (1.89, 2.93)	N/A	4.63 (2.84, 7.54)	N/A	0.84 (0.30, 2.38)	N/A	N/A
<i>P</i>	<0.0001	—	<0.0001	—	0.71	—	—

<sup>1</sup> AUCs are presented as medians (25th, 75th percentiles), *n* = 11 (5 females, 6 males). N/A, not applicable; ND, not determined.

<sup>2</sup> Between tomato sauce co-consumed with avocado vs. sauce alone based on log values presented as geometric means (95% CIs).

A absorption and efficient conversion to vitamin A. In both studies 1 and 2, the bioavailability of provitamin A carotenoids was considerably improved when the test food was consumed with lipid-rich avocado. These results further support previous findings from our group and others (13–16) that increasing amounts of meal lipid increases carotenoid absorption compared with lower amounts of lipid or no lipid. Likewise, a previous study (15) demonstrated that lipid-rich avocado is just as effective as an equivalent amount of pure avocado oil in enhancing carotenoid absorption. When compared with these previous studies (14,15), we observed a smaller magnitude of AUC carotenoid increase when our test meals were consumed with avocado. This difference may be attributed to the larger dose of carotenoid delivered from the sauce in study 1 and from the carrots in study 2 compared with the previous work (11.5 mg of  $\beta$ -carotene and 6.6 mg of  $\alpha$ -carotene) (15). In addition, at higher doses, transporter-facilitated carotenoid absorption was shown to be saturable (32), and, furthermore, carotenoids may compete for absorption (33,34).

In study 2, the ratio of AUC $_{\beta}$ -carotene to AUC $_{\alpha}$ -carotene was almost equal to the ratio of  $\beta$ -carotene to  $\alpha$ -carotene in carrots when the meal was fed with lipid-rich avocado. Thus, under these meal conditions,  $\beta$ -carotene and  $\alpha$ -carotene appear to be absorbed equally. In contrast, this ratio was not maintained when carrots were fed alone, although very little carotenoid was absorbed in general after this test meal. Results from previous human studies are mixed. Some studies reported that carrot  $\beta$ -carotene absorption was approximately double that of carrot  $\alpha$ -carotene when compared on an equimolar basis, as measured by blood response (17) or fecal carotenoid excretion (35). In contrast, other studies reported a greater percentage absorption of  $\alpha$ -carotene relative to  $\beta$ -carotene from carrots after both postprandial (28) and chronic (36) consumption studies. Many factors likely contribute to the disparity between these results.

Strikingly, avocado consumption with the test meals in studies 1 and 2 also led to higher absolute amounts of retinyl

esters (i.e., vitamin A) in the TRL fraction. As a consequence of enhanced carotenoid absorption, the presence of more provitamin A to be converted could at least partially explain the increased appearance of retinyl esters. However, co-consumed lipid may also directly affect other variables that affect conversion, as suggested by a few animal studies. In 1 study, Mongolian gerbils were fed a diet containing carrot powder with 10% lipid (*n* = 12) or 30% lipid (*n* = 12) for 2 wk (22). Animals in the 30% lipid group had considerably higher vitamin A concentrations but lower  $\beta$ -carotene concentrations in liver compared with the 10% lipid group, demonstrating higher conversion with a higher amount of dietary lipid (22). A similar study in ferrets compared the effect of 4 wk of consumption of  $\beta$ -carotene with 6%, 13.4%, or 23% lipid (23). A stepwise increase in dietary lipid was correlated with a stepwise increase in hepatic retinyl ester stores, whereas hepatic  $\beta$ -carotene concentrations for 13.4% or 23% lipid were approximately double those of the 6% group (23). Furthermore, higher consumption of unsaturated lipids was shown to enhance the specific activity of BCO1 in rodents, whereas higher consumption of saturated lipids did not substantially increase BCO1 activity (13). Together, these studies suggest that consuming a higher amount of dietary lipid might increase the conversion rate of provitamin A to vitamin A, particularly when unsaturated lipids (like those found in avocado) are consumed. Besides enhanced enzymatic activity, other research has demonstrated that dietary lipids are necessary for chylomicron synthesis in the enterocyte (37). Thus, increased amounts of retinyl esters in the chylomicron fraction may be a product of increased synthesis and release of chylomicrons containing retinyl esters. Regardless of the mechanism(s) involved, increasing vitamin A formation and delivery to the circulatory system by consuming lipid-rich avocados has practical implications for populations in which vitamin A deficiency is prevalent.

Higher mean conversion rates were reported for  $\beta$ -carotene from various preparations of carrots compared with our


**TABLE 4** Study 2: AUC and fold differences of carotenoids and vitamins after consumption of carrots alone or with avocado in healthy participants<sup>1</sup>

	$\beta$ -Carotene	$\alpha$ -Carotene	Retinyl esters	Lutein	Lycopene	$\alpha$ -Tocopherol	Phylloquinone
	<i>nmol-h/L</i>	<i>nmol-h/L</i>	<i>nmol-h/L</i>	<i>nmol-h/L</i>	<i>nmol-h/L</i>	<i>nmol-h/L</i>	<i>nmol-h/L</i>
Carrot alone (AUC)	88 (24, 125)	70 (31, 97)	51 (22, 97)	34 (3.5, 63)	ND	ND	0.5 (0.0, 0.7)
Carrot with avocado (AUC)	366 (276, 460)	260 (170, 313)	327 (234, 490)	39 (7.4, 70)	ND	1.6 (1.0, 2.8)	4.6 (3.9, 10)
Fold difference <sup>2</sup>	6.63 (4.05, 10.9) <sup>3</sup>	4.83 (3.17, 7.35) <sup>3</sup>	12.6 (3.51, 45.4)	0.77 (0.15, 4.03)	N/A	N/A	15.0 (7.19, 31.3)
<i>P</i>	<0.0001	<0.0001	0.0013	0.73	—	—	<0.0001

<sup>1</sup> AUCs are presented as medians (25th, 75th percentiles), *n* = 12 (6 females, 6 males). N/A, not applicable; ND, not determined.

<sup>2</sup> Fold difference between carrot co-consumed with avocado vs. carrot alone based on log values presented as geometric means (95% CIs).

<sup>3</sup> Fold difference at age 28 y based on log values.




**FIGURE 1** Study 1: baseline-corrected plasma TRL concentrations of  $\beta$ -carotene (A) and retinyl esters (B) over 12 h after consumption of the high- $\beta$ -carotene tomato sauce with avocado and the high- $\beta$ -carotene tomato sauce alone in healthy men and women. Plasma TRL concentrations (nanomoles per liter of plasma) are expressed as means  $\pm$  SEMs,  $n = 11$ . TRL, TG-rich lipoprotein.

study (28). A mean conversion efficiency of 44% was observed for carrot puree, 59% for boiled mashed carrots, and 63% for raw chopped carrots (each delivering 18.6 mg of  $\beta$ -carotene co-consumed with 9 g of safflower oil). In contrast, similar conversion rates were reported when 15 mg of  $\beta$ -carotene was fed in pure oil (33–71% conversion) (38) or palm oil (conversion rate of 69–71%) (39). These studies fed lower doses of  $\beta$ -carotene, which may have contributed to the higher efficiency of conversion.


In study 1, those participants with lower conversion efficiency had a more appreciable increase in conversion with lipid-rich avocado than those with higher efficiency of conversion. It is likely that an individual's vitamin A status will have an impact on provitamin A absorption and subsequent conversion. The liver is the central storage organ of vitamin A and regulator of circulating retinol concentrations (40), whereas blood retinol concentrations remain constant over a wide range of intakes unless an individual is severely deficient or has excessively high consumption (41,42). Thus, the most accurate way to assess vitamin A stores in a healthy population is to estimate liver stores with a deuterated-retinol dilution method (43,44). Because we assumed that all of our participants, as healthy adults living in the United States, had adequate or higher hepatic

vitamin A stores, we did not measure this value. We believe it is more likely that differences in percentage conversion between participants could be related to specific polymorphisms in proteins involved in carotenoid uptake, transport, and/or metabolism in the enterocyte, as reviewed recently (45).

In study 2, the weaker relation of conversion efficiency with and without avocado lipid observed in the carrot study



**FIGURE 2** Study 2: baseline-corrected plasma TRL concentrations of  $\beta$ -carotene (A),  $\alpha$ -carotene (B), and retinyl esters (C) over 12 h after consumption of carrot with avocado and carrot alone in healthy men and women. Plasma TRL concentrations (nanomoles per liter of plasma) are expressed as means  $\pm$  SEMs,  $n = 12$ . TRL, TG-rich lipoprotein.



**FIGURE 3** A plot of the percentage conversion of provitamin A into vitamin A (Efficiency conversion  $A_1$ ) in healthy men and women in study 1, following Eq. 1. Each data point represents a single participant ( $n = 11$ ) and plots their percentage conversion efficiency from the tomato sauce meal alone ( $x$  axis) against their percentage conversion efficiency from the tomato sauce meal with avocado ( $y$  axis). The solid line through the origin (slope = 1) represents equivalent conversion for both meals. The linear equation provided in the graph can be further explained as follows: % Vitamin A conversion<sub>(tomato sauce + avocado)</sub> =  $0.64 \times$  [% Vitamin A conversion<sub>(tomato sauce)</sub>] + 18%. % Vit. A, percentage of vitamin A.

compared with the sauce study may be attributed to the fact that the uptake of carotenes from the carrot meal alone was generally very low (Fig. 2). In contrast, the absorption of  $\beta$ -carotene from the sauce meal alone was much higher. In addition, the difference between the results of the sauce and carrot studies could be explained by a number of factors, for example, the contribution of  $\alpha$ -carotene in carrots to the vitamin A pool. Our decision to assume that  $\alpha$ -carotene provides half as much retinyl ester as  $\beta$ -carotene in Eq. 2 is supported by a previous study with Mongolian gerbils (46). However, this is a mathematical calculation and not a measured value. Differences in food matrix effects (cooked tomato sauce vs. raw carrot) may have also affected the conversion efficiency. Future studies are needed to determine the mechanism(s) responsible.

Vitamin A deficiency represents a real problem in the developing world, and a variety of strategies, including food fortification (47–49), supplementation (50,51), genetically modified rice (52), selective plant breeding of crops with higher provitamin A amounts (53,54), and substituting more nutrient-dense versions of currently consumed vegetables (55–57), were used to ameliorate this problem. Similarly, this novel variety of orange tomatoes could be adapted in vitamin A-deficient regions where red tomatoes are traditionally grown and consumed.

Differences in lutein AUC values were not significantly different for study 2 because of the low and comparable lutein content of the 2 test meals (Table 2). We observed a statistically significant increase in phyloquinone amounts in the carrot study when consumed with avocado compared with carrot alone, likely due to the  $\sim 2.4$ -fold higher dose of phyloquinone in the carrot with avocado meal (Table 2). Studies demonstrating substantial AUC changes for phyloquinone were observed when  $\sim 0.4$ – $1$  mg of this nutrient was fed to humans (58–60)

compared with up to  $0.047$  mg fed in our study. Because it was not the primary aim of our study, participants were not required to abstain from consuming phyloquinone-rich foods during the washout, which may also affected the uptake of this nutrient from the test meal.

In conclusion, consuming provitamin A carotenoids with lipid-rich avocado enhances carotenoid absorption in healthy humans. A notably higher concentration of vitamin A was observed in the TRL fraction when the carotene-rich tomato sauce or carrots were fed with lipid-rich avocado compared with no avocado. Furthermore, consuming lipid-rich avocado with provitamin A from a high- $\beta$ -carotene tomato sauce led to higher conversion efficiency to vitamin A in participants with low conversion efficiency. This observation highlights the importance of consuming provitamin A carotenoids with lipid in the meal, especially in vitamin A-deficient populations in which maximum delivery of active vitamin A is desired.

### Acknowledgments

The authors thank Dr. Dennis Pearl for his advice regarding the statistical analysis of the data and the Hass Avocado Board for providing avocados for the feeding study. R.E.K., D.M.F., E.H.H., S.K.C., and S.J.S. designed the research; R.E.K., J.L.C., and R.M.S. conducted the research; R.E.K., R.M.S., and G.S.Y. analyzed the data; R.E.K. wrote the paper; and S.J.S. had primary responsibility for the final content. All authors read and approved the final manuscript.

### References

1. World Health Organization (WHO). Global prevalence of vitamin A deficiency. Micronutrient Deficiency Information System (MDIS) Working Paper 2 [on the Internet]. Geneva: WHO. 1995 [cited 2012 Sept 13]. Available from: [http://whqlibdoc.who.int/publications/2009/9789241598019\\_eng.pdf](http://whqlibdoc.who.int/publications/2009/9789241598019_eng.pdf).
2. West KP Jr. Vitamin A deficiency disorders in children and women. *Food Nutr Bull* 2003;24:S78–90.
3. Underwood BA, Arthur P. The contribution of vitamin A to public health. *FASEB J* 1996;10:1040–8.
4. Wu J, Taylor RN, Sidell N. Retinoic acid regulates gap junction intercellular communication in human endometrial stromal cells through modulation of the phosphorylation status of connexin 43. *J Cell Physiol* 2013;228:903–10.
5. Lamb TD, Pugh EN. Dark adaptation and the retinoid cycle of vision. *Prog Retin Eye Res* 2004;23:307–80.
6. Durrant A, Harrison E. Digestion and intestinal absorption of dietary carotenoids and vitamin A. *Physiology of the gastrointestinal tract*. 4th ed. Waltham (MA): Academic Press; 2006. p. 1735–52.
7. Calloway DH. Human nutrition: food and micronutrient relationships. Washington: International Food Policy Research Institute; 1995.
8. Neumann CG, Bwibo NO, Murphy SP, Sigman M, Whaley S, Allen LH, Guthrie D, Weiss RE, Demment MW. Animal source foods improve dietary quality, micronutrient status, growth and cognitive function in Kenyan school children: background, study design and baseline findings. *J Nutr* 2003;133:3941S–9S.
9. Zeitlin MF, Megawangi R, Kramer EM, Armstrong HC. Mothers' and children's intakes of vitamin A in rural Bangladesh. *Am J Clin Nutr* 1992;56:136–47.
10. Ramakrishnan U, Martorell R, Latham MC, Abel R. Dietary vitamin A intakes of preschool-age children in South India. *J Nutr* 1999;129:2021–7.
11. von Lintig J, Vogt K. Filling the gap in vitamin A research: molecular identification of an enzyme cleaving  $\beta$ -carotene to retinal. *J Biol Chem* 2000;275:11915–20.
12. Lindqvist A, Andersson S. Cell type-specific expression of  $\beta$ -carotene 15,15'-mono-oxygenase in human tissues. *J Histochem Cytochem* 2004;52:491–9.

13. Roels OA, Trout M, Dujacquier R. Carotene balances in boys in Ruanda where vitamin A deficiency is prevalent. *J Nutr* 1958;65:115–27.
14. Brown MJ, Ferruzzi MG, Nguyen ML, Cooper DA, Eldridge AL, Schwartz SJ, White WS. Carotenoid bioavailability is higher from salads ingested with full-fat than with fat-reduced salad dressings as measured with electrochemical detection. *Am J Clin Nutr* 2004;80:396–403.
15. Unlu NZ, Bohn T, Clinton SK, Schwartz SJ. Carotenoid absorption from salad and salsa by humans is enhanced by the addition of avocado or avocado oil. *J Nutr* 2005;135:431–6.
16. Goltz SR, Campbell WW, Chitchumroonchokchai C, Failla ML, Ferruzzi MG. Meal triacylglycerol profile modulates postprandial absorption of carotenoids in humans. *Mol Nutr Food Res* 2012;56:866–77.
17. Tang G, Qin J, Dolnikowski GG, Russell RM, Grusak MA. Spinach or carrots can supply significant amounts of vitamin A as assessed by feeding with intrinsically deuterated vegetables. *Am J Clin Nutr* 2005;82:821–8.
18. Wang J, Wang Y, Wang Z, Li L, Qin J, Lai W, Fu Y, Suter PM, Russell RM, Grusak MA, et al. Vitamin A equivalence of spirulina  $\beta$ -carotene in Chinese adults as assessed by using a stable-isotope reference method. *Am J Clin Nutr* 2008;87:1730–7.
19. Tang G, Qin J, Dolnikowski GG, Russell RM, Grusak MA. Golden Rice is an effective source of vitamin A. *Am J Clin Nutr* 2009;89:1776–83.
20. Borel P, Tyssandier V, Mekki N, Grolier P, Rochette Y, Alexandre-Gouabau MC, Lairon D, Azaïs-Braesco V. Chylomicron  $\beta$ -carotene and retinyl palmitate responses are dramatically diminished when men ingest  $\beta$ -carotene with medium-chain rather than long-chain triglycerides. *J Nutr* 1998;128:1361–7.
21. During A, Nagao A, Terao J.  $\beta$ -carotene 15,15'-dioxygenase activity and cellular retinol-binding protein type II level are enhanced by dietary unsaturated triacylglycerols in rat intestines. *J Nutr* 1998;128:1614–9.
22. Deming DM, Boileau AC, Lee CM, Erdman JW Jr. Amount of dietary fat and type of soluble fiber independently modulate postabsorptive conversion of  $\beta$ -carotene to vitamin A in Mongolian gerbils. *J Nutr* 2000;130:2789–96.
23. Lakshman MR, Liu QH, Sapp R, Somanchi M, Sundaresan PR. The effects of dietary taurocholate, fat, protein, and carbohydrate on the distribution and fate of dietary beta-carotene in ferrets. *Nutr Cancer* 1996;26:49–61.
24. Unlu NZ, Bohn T, Francis D, Clinton SK, Schwartz SJ. Carotenoid absorption in humans consuming tomato sauces obtained from tangerine or high- $\beta$ -carotene varieties of tomatoes. *J Agric Food Chem* 2007;55:1597–603.
25. Lu QY, Zhang Y, Wang Y, Wang D, Lee R, Gao K, Byrns R, Heber D. California Hass avocado: profiling of carotenoids, tocopherol, fatty acid, and fat content during maturation and from different growing areas. *J Agric Food Chem* 2009;57:10408–13.
26. Kopec RE, Schweiggert RM, Riedl KM, Carle R, Schwartz SJ. Comparison of high-performance liquid chromatography/tandem mass spectrometry and high-performance liquid chromatography/photo-diode array detection for the quantitation of carotenoids, retinyl esters,  $\alpha$ -tocopherol and phyloquinone in chylomicron-rich fractions of human plasma. *Rapid Commun Mass Spectrom* 2013;27:1393–402.
27. Fleshman MK, Riedl KM, Novotny JA, Schwartz SJ, Harrison EH. An LC/MS method for d8- $\beta$ -carotene and d4-retinyl esters:  $\beta$ -carotene absorption and its conversion to vitamin A in humans. *J Lipid Res* 2012;53:820–7.
28. Edwards AJ, Nguyen CH, You CS, Swanson JE, Emenhiser C, Parker RS. Alpha- and beta-carotene from a commercial carrot puree are more bioavailable to humans than from boiled-mashed carrots, as determined using an extrinsic stable isotope reference method. *J Nutr* 2002;132:159–67.
29. Jones B, Kenward MG. Design and analysis of cross-over trials. Boca Raton (FL): Chapman and Hall; 2003.
30. Johnson EJ, Russell RM. Distribution of orally administered  $\beta$ -carotene among lipoproteins in healthy men. *Am J Clin Nutr* 1992;56:128–35.
31. Sies H, Stahl W. Vitamins E and C,  $\beta$ -carotene, and other carotenoids as antioxidants. *Am J Clin Nutr* 1995;62:1315S–21S.
32. During A, Hussain MM, Morel DW, Harrison EH. Carotenoid uptake and secretion by CaCo-2 cells:  $\beta$ -carotene isomer selectivity and carotenoid interactions. *J Lipid Res* 2002;43:1086–95.
33. van het Hof KH, West CE, Weststrate JA, Hautvast JGAJ. Dietary factors that affect the bioavailability of carotenoids. *J Nutr* 2000;130:503–6.
34. Tyssandier V, Cardinault N, Caris-Veyrat C, Amiot MJ, Grolier P, Bouteloup C, Azais-Braesco V, Borel P. Vegetable-borne lutein, lycopene, and  $\beta$ -carotene compete for incorporation into chylomicrons, with no adverse effect on the medium-term (3-wk) plasma status of carotenoids in humans. *Am J Clin Nutr* 2002;75:526–34.
35. Rao CN, Rao BS. Absorption of dietary carotenes in human subjects. *Am J Clin Nutr* 1970;23:105–9.
36. Micozzi MS, Brown ED, Edwards BK, Bieri JG, Taylor PR, Khachik F, Beecher GR, Smith JC. Plasma carotenoid response to chronic intake of selected foods and beta-carotene supplements in men. *Am J Clin Nutr* 1992;55:1120–5.
37. Kalogeris TJ, Monroe F, Demichele SJ, Tso P. Intestinal synthesis and lymphatic secretion of apolipoprotein A-IV vary with chain length of intestinally infused fatty acids in rats. *J Nutr* 1996;126:2720–9.
38. van Vliet T, Schreurs WH, van den Berg H. Intestinal  $\beta$ -carotene absorption and cleavage in men: response of  $\beta$ -carotene and retinyl esters in the triglyceride-rich lipoprotein fraction after a single oral dose of  $\beta$ -carotene. *Am J Clin Nutr* 1995;62:110–6.
39. van den Berg H, van Vliet T. Effect of simultaneous, single oral doses of  $\beta$ -carotene with lutein or lycopene on the  $\beta$ -carotene and retinyl ester responses in the triacylglycerol-rich lipoprotein fraction of men. *Am J Clin Nutr* 1998;68:82–9.
40. Raica N, Jr., Scott J, Lowry L, Sauberlich HE. Vitamin A concentration in human tissues collected from five areas in the United States. *Am J Clin Nutr* 1972;25:291–6.
41. Tanumihardjo SA. Assessing vitamin A status: past, present and future. *J Nutr* 2004;134:290S–3S.
42. Tang G. Bioconversion of dietary provitamin A carotenoids to vitamin A in humans. *Am J Clin Nutr* 2010;91:1468S–73S.
43. Furr HC, Amedee-Manesme O, Clifford AJ, Bergen HR 3rd, Jones AD, Anderson DP, Olson JA. Vitamin A concentrations in liver determined by isotope dilution assay with tetradeuterated vitamin A and by biopsy in generally healthy adult humans. *Am J Clin Nutr* 1989;49:713–6.
44. Ribaya-Mercado JD, Solon FS, Dallal GE, Solomons NW, Fermin LS, Mazariegos M, Dolnikowski GG, Russell RM. Quantitative assessment of total body stores of vitamin A in adults with the use of a 3-d deuterated-retinol-dilution procedure. *Am J Clin Nutr* 2003;77:694–9.
45. Borel P. Genetic variations involved in interindividual variability in carotenoid status. *Mol Nutr Food Res* 2012;56:228–40.
46. Tanumihardjo SA, Howe JA. Twice the amount of  $\alpha$ -carotene isolated from carrots is as effective as  $\beta$ -carotene in maintaining the vitamin A status of Mongolian gerbils. *J Nutr* 2005;135:2622–6.
47. Arroyave G, Mejía LA, Aguilar JR. The effect of vitamin A fortification of sugar on the serum vitamin A levels of preschool Guatemalan children: a longitudinal evaluation. *Am J Clin Nutr* 1981;34:41–9.
48. Dary O, Mora JO. Food fortification to reduce vitamin A deficiency: International Vitamin A Consultative Group recommendations. *J Nutr* 2002;132:2927S–33S.
49. Solon FS, Klemm RD, Sanchez L, Darnton-Hill I, Craft NE, Christian P, West KP. Efficacy of a vitamin A-fortified wheat-flour bun on the vitamin A status of Filipino schoolchildren. *Am J Clin Nutr* 2000;72:738–44.
50. West KP Jr, Pokhrel RP, Katz J, LeClerq SC, Khattry SK, Shrestha SR, Pradhan EK, Tielsch JM, Pandey MR, Sommer A. Efficacy of vitamin A in reducing preschool child mortality in Nepal. *Lancet* 1991;338:67–71.
51. West KP, Jr., Katz J, Khattry SK, LeClerq SC, Pradhan EK, Shrestha SR, Connor PB, Dali SM, Christian P, Pokhrel RP, et al. Double blind, cluster randomised trial of low dose supplementation with vitamin A or  $\beta$ -carotene on mortality related to pregnancy in Nepal. The NNIPS-2 Study Group. *BMJ* 1999;318:570–5.
52. Ye X, Al-Babili S, Klöti A, Zhang J, Lucca P, Beyer P, Potrykus I. Engineering the provitamin A ( $\beta$ -carotene) biosynthetic pathway into (carotenoid-free) rice endosperm. *Science* 2000;287:303–5.
53. Thakkar SK, Maziya-Dixon B, Dixon AGO, Failla ML.  $\beta$ -Carotene micellarization during in vitro digestion and uptake by Caco-2 cells is directly proportional to  $\beta$ -carotene content in different genotypes of cassava. *J Nutr* 2007;137:2229–33.
54. Li S, Nugroho A, Rocheford T, White WS. Vitamin A equivalence of the  $\beta$ -carotene in  $\beta$ -carotene-biofortified maize porridge consumed by women. *Am J Clin Nutr* 2010;92:1105–12.

55. Hagenimana V, Oyunga MA, Low J, Njoroge SM, Gichuki ST, Tuvemo T, Gustafsson J, Proos LA, Worby P, Pitayanon S. The effects of women farmers adoption of orange-fleshed sweet potatoes: raising vitamin A intake in Kenya. *Acta Paediatr* 1999;88:928–32.
56. van Jaarsveld PJ, Faber M, Tanumihardjo SA, Nestel P, Lombard CJ, Benadé AJS.  $\beta$ -Carotene-rich orange-fleshed sweet potato improves the vitamin A status of primary school children assessed with the modified-relative-dose-response test. *Am J Clin Nutr* 2005;81:1080–7.
57. Low JW, Arimond M, Osman N, Cunguara B, Zano F, Tschirley D. A food-based approach introducing orange-fleshed sweet potatoes increased vitamin A intake and serum retinol concentrations in young children in rural Mozambique. *J Nutr* 2007;137:1320–7.
58. Gijssbers BL, Jie KS, Vermeer C. Effect of food composition on vitamin K absorption in human volunteers. *Br J Nutr* 1996;76:223–9.
59. Garber AK, Binkley NC, Krueger DC, Suttie JW. Comparison of phylloquinone bioavailability from food sources or a supplement in human subjects. *J Nutr* 1999;129:1201–3.
60. Booth SL, Lichtenstein AH, Dallal GE. Phylloquinone absorption from phylloquinone-fortified oil is greater than from a vegetable in younger and older men and women. *J Nutr* 2002;132:2609–12.